ELSEVIER

Toxicology 146 (2000) 237–238

TOXICOLOGY

www.elsevier.com/locate/toxicol

# Contents of volume 146

Prolonged elevation in blood pressure in the unrestrained rat exposed to chlorpyrifos
**Cholinergic stimulation and neurohumoral pressor pathways may be involved**
C.J. Gordon, B.K. Padnos (USA) . . . . . 1–13

Inhibition of gap-junctional-intercellular communication in intact rat liver by nongenotoxic hepatocarcinogens
**Agents examined were phenobarbital, pregnenolone-16α-carbonitrile and Aroclor 1254**
K.L. Kolaja, D.T. Engelken, C.D. Klaassen (USA) . . . . . 15–22

Effects of molybdate and pentachlorophenol on the sulfation of acetaminophen
**Pentachlorophenol inhibits sulfation at low doses of acetaminophen whereas molybdate inhibits more effectively at high doses of acetaminophen**
J.W. Boles, C.D. Klaassen (USA) . . . . . 23–35

β-Carotene and protein oxidation: effects of ascorbic acid and α-tocopherol
**Mixtures gave better protection than single compounds**
P. Zhang, S.T. Omaye (USA) . . . . . 37–47

A quantitative method for assessing the sensitizing potency of low molecular weight chemicals using a local lymph node assay: employment of a regression method that includes determination of the uncertainty margins
**A method to quantify the potency of contact sensitizers is proposed**
F.M.M. van Och, W. Slob, W.H. de Jong, R.J. Vandebriel, H. van Loveren (The Netherlands) . . . . . 49–59

Varied protocols of cadmium exposure produce different effects on nitric oxide production in macrophages
**The effect of cadimum on NO generation is dependant on the protocol and timing of exposure**
D.C. Ramirez, M.S. Gimenez (Argentina) . . . . . 61–72

The popliteal lymph node response to streptozotocin is under type 1, MHC class-I restricted, $CD8^+$ T-cell control
**$CD8^+$ T cells are pivotal in the PLN response to streptozotocin**
G. Choquet-Kastylevsky, R. Tedone, M.-T. Ducluzeau, J. Kehren, J.-F. Nicolas, J. Descotes (France). . . . 73–82

Induction of 8-oxo-dGTPase activity in human lymphoid cells and normal fibroblasts by oxidative stress
F. Meyer, E. Fiala, J. Westendorf (Germany, USA) . . . . . 83–92
**Chronic lead application dose-dependently lowers the values for the investigated parameters**

Alterations in β-adrenergic receptor density and adenylate cyclase activity in the rat brain treated chronically with lead
D.-A. Tsao, H.-S. Yu, J.-T. Cheng, C.-K. Ho, H.-R. Chang (Taiwan) . . . . . 93–99

The Long Evans Cinnamon (LEC) rat develops hepatocellular damage in the absence of antimicrosomal antibodies
K. Bedard, I.C. Fuentealba, A. Cribb (Canada) . . . . . 101–109
**Autoantibodies appear to play no role in pathogenesis of hepatitis**

Enhanced expression of rat hepatic microsomal epoxide hydrolase by methylthiazole in conjunction with liver injury
M.K. Cho, S.G. Kim (South Korea) . . . . . 111–122

Trichloroethylene induced vitamin $B_{12}$ and folate deficiency leads to increased formic acid excretion in the rat
J.L. Dow, T. Green (UK) . . . . . 123–136

Developmental lead exposure disturbs expression of synaptic neural cell adhesion molecules in herring gull brains
P.M. Dey, J. Burger, M. Gochfeld, K.R. Reuhl (USA) . . . . . 137–147
**Disruptions in expression of cell adhesive molecules may be a marker for heavy metal exposure during postnatal development**

Interlaboratory comparison of the CB6F1-Tg rasH2 rapid carcinogenicity testing model
R.R. Maronpot, K. Mitsumori, P. Mann, M. Takaoka, S. Yamamoto, T. Usui, H. Okamiya, S. Nishikawa, T. Nomura (USA, Japan) . . . . . 149–159
**Good concordance in outcome for strong carcinogens and non-carcinogens was found**

Effect of long term fluoride exposure on lipid composition in rat liver
Y.-N. Wang, K.-Q. Xiao, J.-L. Liu, G. Dallner, Z.-Z. Guan (PR China, Sweden) . . . . . 161–169
In vitro inhibitory effects of antioxidants on cytotoxicity of T-2 toxin
F. Shokri, M. Heidari, S. Gharagozloo, M. Ghazi-Khansari (Iran) . . . . . 171–176
Influence of microcystine–YR and nodularin on the activity of some glucosidases in mouse liver
A. Lankoff, A. Kołątaj (Poland) . . . . . 177–185
Carrier-mediated uptake of cisplatin by the OK renal epithelial cell line
T. Endo, O. Kimura, M. Sakata (Japan) . . . . . 187–195
**Accumulation of cispolatin into the cell was faster from the basolateral side than from the reverse direction**
Dietary intake of rapeseed oil or soybean oil as the only fat nutrient in spontaneously hypertensive rats and Wistar Kyoto rats — blood pressure and pathophysiology
Y. Naito, H. Yoshida, T. Nagata, A. Tanaka, H. Ono, N. Ohara (Japan) . . . . . 197–208
**Rapeseed oil as only fat source increasers blood pressure in rats under NaCl loading**

**Immunotoxicology articles**
The by-products generated during sarin synthesis in the Tokyo sarin disaster induced inhibition of natural killer and cytotoxic T lymphocyte activity
Q. Li, Y. Hirata, S. Piao, M. Minami (Japan) . . . . . 209–220
**Sarin and its by-products were found to inhibit natural killer and cytotoxic T lymphocyte activity in mice**
Comparison of mouse strains using the local lymph node assay
M.R. Woolhiser, A.E. Munson, B.J. Meade (USA) . . . . . 221–227
**DBA/2, B6c3fl and Balb/c mice are good additional strais for the LLNA**

**News and views** . . . . . 229

**Author index of volume 146** . . . . . 231
**Subject index of volume 146** . . . . . 233–235
**Contents of volume 146** . . . . . 237–238